Brain Research, 440 (1988) 407–409
Elsevier

Author Index

Adinolfi, A.M., see Levine, M.S., 267
Aitken, P.G., see Okazaki, M.M., 352
Allen, S.J., see Dawbarn, D., 185
Amir, S., Aging blocks the thermoregulatory action of thyrotropin-releasing hormone in anaesthetized rats, 181
Awwiller, D.M., see Nordlander, R.H., 391

Bachelard, H.S., see Cox, D.W.G., 195
Bannerman, P.G.C., Mirsky, R. and Jessen, K.R., Antigenic markers and laminin expression in cultured enteric neural cells, 87
Bannerman, P.G.C., Mirsky, R. and Jessen, K.R., Establishment and properties of separate cultures of enteric neurons and enteric glia, 99
Baranyi, A., Szente, M.B. and Woody, C.D., Activation of protein kinase C induces long-term changes of postsynaptic currents in neocortical neurons, 341
Barberis, C., see Petit, C.M., 299
Baron, P., see Scarpini, E., 261
Barrantes, F.J., see Ginobili de Martinez, M.S., 386
Bastian, F.O., see Schlitt, M., 293
Beretta, S., see Scarpini, E., 261
Berge, O.-G., see Eide, P.K., 42
Bhargava, H.N., Matwyshyn, G.A., Hanissian, S. and Tejwani, G.A., Opioid peptides in pituitary gland, brain regions and peripheral tissues of spontaneously hypertensive and Wistar–Kyoto normotensive rats, 333
Blanc, G., see Taghzouti, K., 172
Bland, B.H. and Colom, L.V., Responses of phasic and tonic hippocampal theta-on cells to cholinergics: differential effects of muscarinic and nicotinic activation, 167
Boman, A., see Christenson, J., 1
Breitbach, D., see Hayward, L., 199
Brightman, M.W., see Nagy, Z., 315
Broch, O.J., see Eide, P.K., 42
Bucher, A.P., see Schlitt, M., 293
Buchwald, N.A., see Levine, M.S., 267
Burde, R.M., Disparate visceral neuronal pools subserve spinal cord and ciliary ganglion in the monkey: a double labeling approach, 177
Burton, N.R. and Williams, T.L., Fictive locomotion in the lamprey; a comparison using bicarbonate and HEPES buffers, 159
Buscaglia, M., see Scarpini, E., 261

Calaresu, F.R., see Ciriello, J., 53
Camhi, J.M., see Goldstein, R.S., 366
Camxs, M., see Closse, A., 123
Cannon-Spoor, E., see Karoum, F., 190
Caverson, M.M., see Ciriello, J., 53
Choi, D.W., see George, C.P., 375
Chou, Y-.C. and Luttge, W.G., Activated Type II receptors in brain cannot rebind glucocorticoids: relationship to progesterone's antiglucocorticoid actions, 67
Christenson, J., Boman, A., Lagerbäck, P.-Å. and Grillner, S., The dorsal cell, one class of primary sensory neuron in the lamprey spinal cord. I. Touch, pressure but no nociception — a physiological study, 1
Christenson, J., Lagerbäck, P.Å. and Grillner, S., The dorsal cell, one class of primary sensory neuron in the lamprey spinal cord. II. A light- and electron microscopical study, 9
Chuang, L.-W., see Karoum, F., 190
Ciriello, J., Caverson, M.M., Calaresu, F.R. and Krukoff, T.L., Neuropeptide and serotonin immunoreactive neurons in the cat ventrolateral medulla, 53
Closse, A., Camps, M., Wanner, A. and Palacios, J.M., In vivo labeling of brain dopamine D_2 receptors using the high-affinity specific D_2 agonist [^{3}H]CV 205-502, 123
Colom, L.V., see Bland, B.H., 167
Cook, H., see Nordlander, R.H., 391
Costa, E., see Karoum, F., 190
Cox, D.W.G. and Bachelard, H.S., On the relationship between the excitability of dentate granule cell field potentials and their sensitivity to low glucose, 195

Dawbarn, D., Allen, S.J. and Semenenko, F.M., Immunohistochemical localization of β-nerve growth factor receptors in the forebrain of the rat, 185
De Caro, G., see Massi, M., 232
Donnelly, D.F., see Sica, A.L., 370
Dragunow, M. and Robertson, H.A., Localization and induction of c-fos protein-like immunoreactive material in the nuclei of adult mammalian neurons, 252
Dykes, R. and Lamour, Y., Neurons without demonstrable receptive fields outnumber neurons having receptive fields in samples from the somatosensory cortex of anesthetized or paralyzed cats and rats, 133

Eide, P.K., Hole, K., Berge, O.-G. and Broch, O.J., 5-HT depletion with 5,7-DHT, PCA and PCPA in mice: differential effects on the sensitivity to 5-MeODMT, 8-OH-DPAT and 5-HTP as measured by two nociceptive tests, 42
Epstein, A.N., see Massi, M., 232
Eysel, U.T., see Pape, H.-C., 79

Fisher, R.S., see Levine, M.S., 267
Freed, W.J., see Karoum, F., 190
Frey, U., see Reymann, K.G., 305

Gandhi, M.R., see Sica, A.L., 370
Garthwaite, G. and Garthwaite J., Electron microscopic autoradiography of D-[^{3}H]aspartate uptake sites in mouse cerebellar slices shows poor labelling of mossy fibre terminals, 162
Garthwaite, J., see Garthwaite G., 162
Gebhart, G.F., see Janss, A.J., 109
George, C.P., Goldberg, M.P., Choi, D.W. and Steinberg, G.K., Dextromethorphan reduces neocortical ischemic neuronal damage in vivo, 375
Ginobili de Martinez, M.S. and Barrantes, F.J., Ca^{2+} and phospholipid-dependent protein kinase activity in rat cerebral hemispheres, 386
Gioannini, T.L., see Keren, O., 280
Glowinski, J., see Taghzouti, K., 172

Goldberg, M.P., see George, C.P., 375
Goldstein, R.S. and Camhi, J.M., Reduction of sensory activity produced by cercal displacement modifies response of wind-sensitive interneurons in the cockroach, 366
Graczak, L.M., see Quock, R.M., 35
Grillner, S., see Christenson, J., 1
Grillner, S., see Christenson, J., 9
Guthrie, D., see Levine, M.S., 267

Hanissian, S., see Bhargava, H.N., 333
Hasegawa, H., see Kakudo, K., 153
Hashimoto, P.H., Tracer in cisternal cerebrospinal fluid is soon detected in choroid plexus capillaries, 149
Hayward, L., Breitbach, D. and Rymer, W.Z., Increased inhibitory effects on close synergists during muscle fatigue in the decerebrate cat, 199
Hervé, D., see Taghzouti, K., 172
Hiller, J.M., see Keren, O., 280
Hole, K., see Eide, P.K., 42
Hull, C.D., see Levine, M.S., 267

Itoh, Y., see Kakudo, K., 153

Janss, A.J. and Gebhart, G.F., Brainstem and spinal pathways mediating descending inhibition from the medullary lateral reticular nucleus in the rat, 109
Jard, S., see Petit, P., 299
Jennings, R.A., see Schlitt, M., 293
Jessen, K.R., see Bannerman, P.G.C., 87
Jessen, K.R., see Bannerman, P.G.C., 99
Joëls, M., see Schiess, M.C., 380
Jork, R., see Reymann, K.G., 305

Kakudo, K., Hasegawa, H., Komatsu, N., Nakamura, A., Itoh, Y. and Watanabe, K., Immuno-electron microscopic study of calcitonin gene-related peptide (CGRP) in axis cylinders of the vagus nerve. CGRP is present in both myelinated and unmyelinated fibers, 153
Kalin, N.H., Shelton, S.E. and Barksdale, C.M., Opiate modulation of separation-induced distress in non-human primates, 285
Kamphuis, W., Lopes da Silva, F.H. and Wadman, W.J., Changes in local evoked potentials in the rat hippocampus (CA_1) during kindling epileptogenesis, 205
Karoum, F., Freed, W.J., Chuang, L.-W., Cannon-Spoor, E., Wyatt, R.J. and Costa, E., D-DOPA and L-DOPA similarly elevate brain dopamine and produce turning behavior in rats, 190
Keizer, R.F., see Kim, C.S., 216
Kendrick, K.M. and Leng, G., Haemorrhage-induced release of noradrenaline, 5-hydroxytryptamine and uric acid in the supraoptic nucleus of the rat, measured by microdialysis, 402
Keren, O., Gioannini, T.L., Hiller, J.M. and Simon, E.J., Affinity crosslinking of ^{125}I-labeled human β-endorphin to cell lines possessing either μ- or δ-type opioid binding sites, 280
Kim, C.S., Keizer, R.F. and Pritchard, J.B., 2,4-Dichlorophenoxyacetic acid intoxication increases its accumulation within the brain, 216
Koh, S. and Loy, R., Age-related loss of nerve growth factor sensitivity in rat basal forebrain neurons, 396
Komatsu, N., see Kakudo, K., 153
Kreider, B.Q., see Scarpini, E., 261
Krukoff, T.L., see Ciriello, J., 53

Lagerbäck, P.-Å., see Christenson, J., 1
Lagerbäck, P.-Å., see Christenson, J., 9
Lakeman, A.D., see Schlitt, M., 293
Lamour, Y., see Dykes, R., 133
LeMoal, M., see Taghzouti, K., 172
Leng, G., see Kendrick, K.M., 402
Letourneau, P.C., see Quick, D.C., 243
Levine, M.S., Adinolfi, A.M., Fisher, R.S., Hull, C.D., Guthrie, D. and Buchwald, N.A., Ultrastructural alterations in caudate nucleus in aged cats, 267
Lisak, R.P., see Scarpini, E., 261
Lopes da Silva, F.H., see Kamphuis, W., 205
Loy, R., see Koh, S., 396
Luttge, W.G., see Chou, Y.-C., 67

Mackel, R. and Noda, T., Sensory input to cerebellocerebral relay neurons in the cat thalamus, 348
Massi, M., Perfumi, M., De Caro, G. and Epstein, A.N., Inhibitory effect of kassinin on salt intake induced by different natriorexigenic treatments in the rat, 232
Mato, M., see Nishimura, Y., 144
Matsushima, R., see Tashiro, T., 357
Matthies, H., see Reymann, K.G., 305
Matwyshyn, G.A., see Bhargava, H.N., 333
Meiselman, S., see Nagy, Z., 315
Meola, G., see Scarpini, E., 261
Mirsky, R., see Bannerman, P.G.C., 87
Mirsky, R., see Bannerman, P.G.C., 99
Mizuno, N., see Tashiro, T., 357
Mooradian, A.D., Blood–brain barrier transport of choline is reduced in the aged rat, 328
Morello, H. and Taleisnik, S., The inhibition of proestrous LH surge and ovulation in rats bearing lesions of the dorsal raphe nucleus is mediated by the locus coeruleus, 227
Moretto, A. and Sabri, M.I., Progressive deficits in retrograde axon transport precede degeneration of motor axons in acrylamide neuropathy, 18

Nadler, J.V., see Okazaki, M.M., 352
Nagy, Z., Pettigrew, K.D., Meiselman, S. and Brightman, M.W., Cerebral vessels cryofixed after hyperosmosis or cold injury in normothermic, and hypothermic frogs, 315
Nakamura, A., see Kakudo, K., 153
Natori, M., see Nishimura, Y., 144
Nishimura, Y., Natori, M. and Mato, M., Choline acetyltransferase immunopositive pyramidal neurons in the rat frontal cortex, 144
Noda, T., see Mackel, R., 348
Nordlander, R.H., Awwiller, D.M. and Cook, H., Dorsal roots are absent from the tail of larval *Xenopus*, 391

Okazaki, M.M., Aitken, P.G. and Nadler, J.V., Mossy fiber lesion reduces the probability that kainic acid will provoke CA3 hippocampal pyramidal cell bursting, 352
Orrego, F., see Villanueva, S., 363

Palacios, J.M., see Closse, A., 123
Pape, H.-C. and Eysel, U.T., Cholinergic excitation and inhibition in the visual thalamus of the cat — influences of cortical inactivation and barbiturate anesthesia, 79
Perfumi, M., see Massi, M., 232
Petit, P., Barberis, C. and Jard, S., Vasopressin potentiates the noradrenaline-induced accumulation of cyclic AMP in the rat superior cervical ganglion, 299

Pettigrew, K.D., see Nagy, Z., 315
Pindak, F., see Schlitt, M., 293
Prasad, N., see Sica, A.L., 370
Pritchard, J.B., see Kim, C.S., 216

Quick, D.C. and Letourneau, P.C., Immuno-labelling for freeze-fracture: application to a cell surface attachment antigen, 243
Quock, R.M. and Graczak, L.M., Influence of narcotic antagonist drugs upon nitrous oxide analgesia in mice, 35

Reymann, K.G., Frey, U., Jork, R. and Matthies, H., Polymyxin B, an inhibitor of protein kinase C, prevents the maintenance of synaptic long-term potentiation in hippocampal CA_1 neurons, 305
Robertson, H.A., see Dragunow, M., 252
Ross, A.H., see Scarpini, E., 261
Rymer, W.Z., see Hayward, L., 199

Sabri, M.I., see Moretto, A., 18
Satoda, T., see Tashiro, T., 357
Scarlato, G., see Scarpini, E., 261
Scarpini, E., Kreider, B.Q., Lisak, R.P., Meola, G., Velicogna, M.E., Baron, P., Beretta, S., Buscaglia, M., Ross, A.H. and Scarlato, G., Cultures of human Schwann cells isolated from fetal nerves, 261
Schiess, M.C., Joëls, M. and Shinnick-Gallagher, P., Estrogen priming affects active membrane properties of medial amygdala neurons, 380
Schlitt, M., Bucher, A.P., Stroop, W.G., Pindak, F., Bastian, F.O., Jennings, R.A., Lakeman, A.D. and Whitley, R.J., Neurovirulence in an experimental focal herpes encephalitis: relationship to observed seizures, 293
Schwarting, G.A., see Suchy, S.F., 25
Semenenko, F.M., see Dawbarn, D., 185
Shelton, S.E., see Kalin, N.H., 285
Shinnick-Gallagher, P., see Schiess, M.C., 380
Sica, A.L., Steele, A.M., Gandhi, M.R., Donnelly, D.F. and Prasad, N., Power spectral analyses of inspiratory activities in neonatal pigs, 370
Simon, E.J., see Keren, O., 280
Simon, H., see Taghzouti, K., 172
Steele, A.M., see Sica, A.L., 370
Steinberg, G.K., see George, C.P., 375
Stroop, W.G., see Schlitt, M., 293
Studler, J.M., see Taghzouti, K., 172
Suchy, S.F., Yamamoto, M., Barbero, L. and Schwarting, G.A., A monoclonal antibody, WCC4, recognizes a developmentally regulated ganglioside containing α-galactose and α-fucose present in the rat nervous system, 25
Szente, M.B., see Baranyi, A., 341

Taghzouti, K., Simon, H., Hervé, D., Blanc, G., Studler, J.M., Glowinski, J., LeMoal, M. and Tassin, J.P., Behavioural deficits induced by an electrolytic lesion of the rat ventral mesencephalic tegmentum are corrected by a superimposed lesion of the dorsal noradrenergic system, 172
Takahashi, O., see Tashiro, T., 357
Taleisnik, S., see Morello, H., 227
Tashiro, T., Satoda, T., Takahashi, O., Matsushima, R. and Mizuno, N., Distribution of axons exhibiting both enkephalin- and serotonin-like immunoreactivities in the lumbar cord segments: an immunohistochemical study in the cat, 357
Tassin, J.P., see Taghzouti, K., 172
Tejwani, G.A., see Bhargava, H.N., 333

Velicogna, M.E., see Scarpini, E., 261
Villanueva, S. and Orrego, F., Endogenous ligands for the quisqualate receptor: presence in rat brain cortex synaptic vesicles, 363

Wadman, W.J., see Kamphuis, W., 205
Wanner, A., see Closse, A, 123
Watanabe, K., see Kakudo, K., 153
Whitley, R.J., see Schlitt, M., 293
Williams, T.L., see Burton, N.R., 159
Woody, C.D., see Baranyi, A., 341
Wyatt, R.J., see Karoum, F., 190

Yamamoto, M., see Suchy, S.F., 25